

Petition for *Inter Partes* Review of USP 9,006,224

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re *Inter Partes Review* of:)
U.S. Patent No. 9,006,224)
Issued: Apr. 14, 2015)
Application No.: 12/094,173)
PCT Filing Date: Nov. 20, 2006)

For: **Neuroendocrine Tumor Treatment**

FILED VIA E2E

**PETITION FOR *INTER PARTES* REVIEW
OF U.S. PATENT NO. 9,006,224**

TABLE OF CONTENTS

TABLE OF AUTHORITIES iv

I. OVERVIEW 1

II. REQUIREMENTS FOR PETITION FOR INTER PARTES REVIEW 2

 A. Grounds for Standing (37 C.F.R. § 42.104(a)) 2

 B. Notice of Lead and Backup Counsel and Service Information..... 3

 C. Notice of Real-Parties-in-Interest (37 C.F.R. § 42.8(b)(1))..... 3

 D. Notice of Related Matters (37 C.F.R. § 42.8(b)(2))..... 4

 E. Fee for *Inter Partes* Review 5

 F. Proof of Service..... 5

III. IDENTIFICATION OF CLAIMS BEING CHALLENGED (37 C.F.R. § 42.104(B)) 6

IV. SUMMARY OF THE ARGUMENT 6

V. OVERVIEW OF THE '224 PATENT 15

VI. THE PERSON OF ORDINARY SKILL IN THE ART 18

VII. CLAIM CONSTRUCTION 19

 A. Applicable Law 19

 B. Construction of Claim Terms 21

 1. “pancreatic neuroendocrine tumor” 22

 2. “advanced tumors” 23

 3. “unit dose” 24

 4. “islet cell tumor” 25

VIII. TECHNICAL BACKGROUND AND STATE OF THE ART AT THE TIME OF THE PURPORTED INVENTION 26

 A. Rapamycin was well-known as a potent antitumor agent..... 26

 B. Rapamycin derivatives, like everolimus and temsirolimus, were known to have similar biological activity to rapamycin 26

Petition for *Inter Partes* Review of USP 9,006,224

- C. The mechanism of action for the immunosuppressant and antitumor activity of rapamycin and its derivatives was well-characterized..... 31
- IX. THE SCOPE AND CONTENT OF THE ASSERTED PRIOR ART 34
 - A. Öberg 2004 taught that humans with advanced pancreatic NETs should be treated with rapamycin as a monotherapy after cytotoxic therapy failed..... 34
 - B. Boulay 2004 taught that everolimus was well-tolerated and effective at treating pancreatic NETs in rat models 37
 - C. O’Donnell taught that everolimus was well-tolerated and showed promise as an antitumor agent in human patients..... 40
 - D. Tabernero taught that an appropriate dosage for humans taking everolimus for the treatment of advanced solid tumors was 10 mg/day 41
 - E. Duran taught the use of temsirolimus in the treatment of human patients with advanced neuroendocrine carcinomas 42
- X. CLAIMS 1-3 WOULD HAVE BEEN OBVIOUS OVER THE PRIOR ART 43
 - A. Legal Background 43
 - B. Ground 1: Claims 1-3 would have been obvious in view of Öberg 2004, Boulay 2004, and O’Donnell 46
 - 1. Claim 1 46
 - 2. Claim 2 53
 - 3. Claim 3 55
 - C. Ground 2: Claim 2 would have been obvious in view of Öberg 2004, Boulay 2004, O’Donnell, and Tabernero 55
 - D. Ground 3: Claims 1-3 would have been obvious in view of Boulay 2004, O’Donnell, and Duran 56
 - 1. Claim 1 56
 - 2. Claim 2 58
 - 3. Claim 3 60

Petition for *Inter Partes* Review of USP 9,006,224

E. Ground 4: Claim 2 of the '224 patent is invalid as obvious in view of Boulay 2004, O'Donnell, Duran, and Taberero..... 61

XI. SECONDARY CONSIDERATIONS FAIL TO OVERCOME THE STRONG EVIDENCE OF OBVIOUSNESS 62

XII. CONCLUSION..... 64

TABLE OF AUTHORITIES

	Page(s)
Cases	
<i>Bayer Healthcare Pharm., Inc. v. Watson Pharm., Inc.</i> , 713 F.3d 1369 (Fed. Cir. 2013)	63
<i>In re Beattie</i> , 974 F.2d 1309 (Fed. Cir. 1992)	14, 52
<i>Boston Sci. Scimed, Inc. v. Cordis Corp.</i> , 554 F.3d 982 (Fed. Cir. 2009)	46
<i>Cuozzo Speed Techs., LLC v. Lee</i> , 136 S.Ct. 2131,2142 (2016).....	19
<i>Galderma Labs., L.P. v. Tolmar, Inc.</i> , 737 F.3d 731 (Fed. Cir. 2013)	63
<i>Graham v. John Deere Co.</i> , 383 U.S. 1 (1966).....	44, 62
<i>KSR Int’l Co. v. Teleflex Inc.</i> , 550 U.S. 398 (2007).....	<i>passim</i>
<i>Leapfrog Enters., Inc. v. Fisher-Price, Inc.</i> , 485 F.3d 1157 (Fed. Cir. 2007)	62
<i>Merck & Co. v. Teva Pharm. USA, Inc.</i> , 395 F.3d 1364 (Fed. Cir. 2005)	53-54, 59, 63
<i>Microsoft Corp. v. Proxyconn, Inc.</i> , 789 F.3d 1292 (Fed. Cir. 2015)	20
<i>In re PepperBall Techs., Inc.</i> , 469 F. App’x 878 (Fed. Cir. 2012)	63

Petition for *Inter Partes* Review of USP 9,006,224

Pfizer, Inc. v. Apotex, Inc.,
480 F.3d 1348 (Fed. Cir. 2007)54, 60

Phillips v. AWH Corp.,
415 F.3d 1303 (Fed. Cir. 2005) (en banc) 19-20

Ruiz v. A.B. Chance Co.,
234 F.3d 654 (Fed. Cir. 2000)44

SIBIA Neurosciences, Inc. v. Cadus Pharm. Corp.,
225 F.3d 1349 (Fed. Cir. 2000)63

Versata Dev. Grp., Inc. v. SAP Am., Inc.,
793 F.3d 1306 (Fed. Cir. 2015)20

Statutes

35 U.S.C. § 102(a)41, 42

35 U.S.C. § 102(b)34, 37, 40

35 U.S.C. § 1036, 43

35 U.S.C. § 301(a)21

35 U.S.C. § 318(a)2

Other Authorities

37 C.F.R. § 42.8(b)(1).....3

37 C.F.R. § 42.8(b)(2).....4

37 C.F.R. § 42.8(b)(3).....3

37 C.F.R. § 42.8(b)(4).....3

37 C.F.R. § 42.10(a).....3

37 C.F.R. § 42.10(b)3

Petition for *Inter Partes* Review of USP 9,006,224

37 C.F.R. § 42.15(a).....5
37 C.F.R. § 42.100(b)19
37 C.F.R. § 42.104(a).....2
37 C.F.R. § 42.104(b)6
77 Fed. Reg. 48680, 48699 (Aug. 14, 2012)20
77 Fed. Reg. 48759-60.....4

P.T.A.B. Decision

SAP Am., Inc. v. Versata Dev. Grp., Inc.,
No. CBM2012-00001 (P.T.A.B. Jan. 9, 2013).....21

Petition for *Inter Partes* Review of USP 9,006,224

Exhibit List

1001	U.S. Patent No. 9,006,224 (“the ’224 patent”), titled “Neuroendocrine Tumor Treatment”
1002	File History for the ’224 patent
1003	Declaration of Mark J. Ratain, M.D. in Support of Petition for <i>Inter Partes</i> Review of U.S. Patent No. 9,006,224 (“Ratain Decl.”)
1004	Curriculum Vitae of Mark J. Ratain, M.D.
1005	A. Boulay et al., <i>Antitumor efficacy of intermittent treatment schedules with the rapamycin derivative RAD001 correlates with Prolonged Inactivation of Ribosomal Protein S6 Kinase 1 in Peripheral Blood Mononuclear Cells</i> , 64 CANCER RES. 252 (2004) (“Boulay 2004”)
1006	E. Brown et al., <i>A mammalian protein targeted by G1-arresting rapamycin-receptor complex</i> , 369 NATURE 756 (1994) (“Brown”)
1007	P. Buetow et al., <i>Islet cell tumors of the Pancreas: Pathologic-Imaging Correlation Among Size, Necrosis and Cysts, Calcification, Malignant Behavior, and Functional Status</i> , 165 AM. J. ROENTGENOLOGY 1175 (1995)
1008	Center for Drug Evaluation & Research, <i>Approval Package for NDA 021083 (Rapamune)</i> , Food & Drug Administration (Sept. 15, 1999)
1009	J. Dancey, <i>Clinical development of mammalian target of rapamycin inhibitors</i> , 16 HEMATOLOGY/ONCOLOGY CLINICS OF N. AM. 1101 (2002)(“Dancey”)
1010	M. DeJong et al., <i>Therapy of neuroendocrine tumors with radiolabeled somatostatin-analogues</i> , 43 Q. J. NUCLEAR MED. & MOLECULAR IMAGING 356 (1999) (“DeJong”)
1011	I. Duran et al., <i>A Phase II Trial of Temsirolimus in Metastatic Neuroendocrine Carcinomas (NECs)</i> , 23 SUPPL. J. CLIN. ONCOL. 3096 (2005) (“Duran”)

Petition for *Inter Partes* Review of USP 9,006,224

1012	J. Dutcher, <i>Mammalian target of rapamycin inhibition</i> , 10 CLIN. CANCER RES. 6382s (2004) (“Dutcher”)
1013	C. P. Eng et al., <i>Activity of Rapamycin (AY-22,989) Against Transplanted Tumors</i> , 37 J. ANTIBIOTICS 1231 (1984) (“Eng”)
1014	M. Grewe et al., <i>Regulation of Cell Growth and Cyclin D1 Expression by the Constitutively Active FRAP-p70^{s6K} Pathway in Human Pancreatic K Cancer Cells</i> , 59 CANCER RES. 3581 (1999) (“Grewe”)
1015	M. Guba et al., <i>Rapamycin inhibits primary and metastatic tumor growth by antiangiogenesis: involvement of vascular endothelial growth factor</i> , 8 NATURE MED. 128 (2002) (“Guba”)
1016	M. Hidalgo et al., <i>The rapamycin-sensitive signal transduction pathway as a target for cancer therapy</i> , 19 ONCOGENE 6680 (2000) (“Hidalgo”)
1017	S. Huang et al., <i>Inhibitors of mammalian target of rapamycin as novel antitumor agents: from bench to clinic</i> , 3 CURRENT OPINION IN INVESTIGATIONAL DRUGS 295 (2002) (“Huang 2002”)
1018	S. Huang et al., <i>Rapamycins: Mechanism of Action and Cellular Resistance</i> , 2 CANCER BIOL. & THER. 222 (2003) (“Huang 2003”)
1019	M. Levy and M. Wiersema, <i>Pancreatic neoplasms</i> , 15 GASTROINTESTINAL ENDOSCOPY CLIN. N. AM. 117 (2005) (“Levy”)
1020	G. Kaltsas et al., <i>The Diagnosis and Medical Management of Advanced Neuroendocrine Tumors</i> , 25 ENDOCRINE REV. 458 (2004) (“Kaltsas”)
1021	R. Martel et al., <i>Inhibition of the immune response by rapamycin, a new antifungal antibiotic</i> , 55 CAN. J. PHYSIOL. PHARMACOL. 48 (1977) (“Martel”)
1022	R. Morris, <i>Rapamycins: Antifungal, Antitumor, Antiproliferative, and Immunosuppressive Macro/ides</i> , 6 TRANSPLANTATION REV. 39 (1992) (“Morris”)

Petition for *Inter Partes* Review of USP 9,006,224

1023	C. Moertel et al., <i>Streptozocin-Doxorubicin, Streptozocin-Fluorouracil, or Chlorozotocin in the Treatment of Advanced Islet-Cell Carcinoma</i> , 326 NEW ENG. J. MED. 519 (1992) (“Moertel”)
1024	M. Neshat et al., <i>Enhanced sensitivity of PTEN-deficient tumors to inhibition of FRAP/mTOR</i> , 98 PNAS 10314 (2001) (“Neshat”)
1025	K. Öberg, <i>Chemotherapy and biotherapy in the treatment of neuroendocrine tumours</i> , 12 ANN. ONCOL. S111 (2001) (“Öberg 2001”)
1026	K. Öberg, <i>Management of neuroendocrine tumors</i> , 15 ANN. ONCOLOGY iv293 (2004)
1027	K. Öberg, <i>Treatment of neuroendocrine tumors of the gastrointestinal tract</i> , 27 ONCOLOGIA 57 (2004) (“Öberg 2004”)
1028	K. Öberg and B. Eriksson, <i>Endocrine tumours of the pancreas</i> , 19 BEST PRACTICE & RES. CLIN. GASTROENT. 753 (2005) (“Öberg & Eriksson”)
1029	A. O’Donnell et al., <i>A phase I study of the oral mTOR inhibitor RAD001 as a monotherapy to identify the optimal biologically effective dose using toxicity, pharmacokinetic (PK) and pharmacodynamics (PD) endpoints in patients with solid tumors</i> , 22 PROC. AM. SOC’Y OF CLINICAL ONCOLOGY 200(803ab) (2003) (“O’Donnell”)
1030	T. O’Reilly et al., <i>In vivo activity of RAD001, an orally active rapamycin derivative, in experimental tumor models</i> , 43 PROC. AM. ASS’N OF CANCER RES. 71 (Abstract #359) (2002) (“O’Reilly”)
1031	A. Perren, et al., <i>Mutation and expression analyses reveal differential subcellular compartmentalization of PTEN in endocrine pancreatic tumors compared to normal/islet cells</i> , 157 AM. J. PATHOLOGY 1097 (2000) (“Perren”)
1032	U. Plockinger et al., <i>Guidelines for the Diagnosis and Treatment of Neuroendocrine Gastrointestinal Tumours</i> , 80 NEUROENDOCRINOLOGY 394 (2004) (“NET Guidelines”)

Petition for *Inter Partes* Review of USP 9,006,224

1033	R. Rao et al., <i>Mammalian Target of Rapamycin (mTOR) Inhibitors as Anti-Cancer Agents</i> , 4 CURR. CANCER DRUG TARGETS 621 (2004) (“Rao”)
1034	C. Sawyers, <i>Will mTOR inhibitors make it as cancer drugs?</i> , 4 CANCER CELL 343 (2003) (“Sawyers”)
1035	S. Schreiber, <i>Chemistry and biology of the immunophilins and their immunosuppressive ligands</i> , 251 SCIENCE 283 (1991) (“Schreiber”)
1036	W. Schuler et al., <i>SDZ RAD, a new rapamycin derivative: pharmacological properties in vitro and in vivo</i> , 64 TRANSPLANTATION 36 (1997) (“Schuler”)
1037	A. Tolcher, <i>Novel therapeutic molecular targets for prostate cancer: the mTOR signaling pathway and epidermal growth factor receptor</i> , 171 J. UROLOGY S41 (2004) (“Tolcher”)
1038	J. Taberero et al., <i>A phase I study with tumor molecular pharmacodynamic (MPD) evaluation of dose and schedule of the oral mTOR-inhibitor Everolimus (RAD001) inpatients (pts) with advanced solid tumors</i> , 23 J. CLINICAL ONCOLOGY 3007 (2005) (“Taberero”)
1039	S. Vignot et al., <i>mTOR-targeted therapy of cancer with rapamycin derivatives</i> , 16 ANN. ONCOL. 525 (2005) (“Vignot”)
1040	U.S. Patent No. 3,929,992 (“the ’992 patent”)
1041	U.S. Patent No. 4,650,803 (“the ’803 patent”)
1042	U.S. Patent No. 4,885,171 (“the ’171 patent”)
1043	U.S. Patent No. 5,100,883 (“the ’883 patent”)
1044	U.S. Patent No. 5,206,018 (“the ’018 patent”)
1045	U.S. Patent No. 5,233,036 (“the ’036 patent”)
1046	U.S. Patent No. 5,362,718 (“the ’718 patent”)
1047	U.S. Patent No. 5,391,730 (“the ’730 patent”)

Petition for *Inter Partes* Review of USP 9,006,224

1048	U.S. Patent No. 5,665,772 (“the ’772 patent”)
1049	U.S. Patent No. 7,091,213 (“the ’213 patent”)
1050	U.S. Patent No. 8,410,131 (“the ’131 patent”)
1051	L. Wang et al., <i>Differential Expression of the PTEN Tumor Suppressor Protein in Fetal and Adult Neuroendocrine Tissues and Tumors: Progression Loss of PTEN Expression in Poorly Differentiated Neuroendocrine Neoplasms</i> , 10 APPLIED IMMUNOHISTOCHEMISTRY & MOLECULAR MORPHOLOGY 139 (2002) (“Wang 2002”)
1052	B. Wiedenmann & U. Pape, <i>From Basic to Clinical Research in Gastroenteropancreatic Neuroendocrine Tumor Disease-The Clinician-Scientist Perspective</i> , 80 NEUROENDOCRINOLOGY 94 (2004) (“Wiedenmann 2004”)
1053	WO 97/47317 (“Weckbecker”)
1054	WO 02/40000 (“Dukart”)
1055	WO 02/066019 (“Lane”)
1056	Excerpt from the file history of U.S. Application No. 14/608,644, Information Disclosure Statement (April, 2015)
1057	Excerpt from the file history of U.S. Application No. 14/608,644, Office Action (December 18, 2015)
1058	Dr. Kjell Öberg, Web Bio, Uppsala University, available at http://katalog.uu.se/empinfo?languageld=1&id=n96-5147 , visited June 26, 2015 (“Öberg Biography”)
1059	“What is ENETS?,” available at http://www.enets.org/what_is_enets.html , visited June 26, 2015 (“ENETS Info”)
1060	Sandostatin LAR [®] Prescribing Label (November 1998)

Petition for *Inter Partes* Review of USP 9,006,224

1061	Declaration of Scott Bennett, Ph.D.
1062	D. Clements et al., <i>Regression of Metastatic Vipoma with Somatostatin Analogue SMS 201-995</i> , LANCET 874 (1985) (“Clements”)
1063	D. O’Toole et al., <i>Chemotherapy for Gastro-Enteropancreatic Endocrine Tumours</i> 80 NEUROENDOCRINOLOGY 79 (2004)
1064	A. Jimeno et al., <i>Pharmacodynamic-guided, modified continuous reassessment method (mCRM)-based, dose finding study of rapamycin in adult patients with solid tumors</i> , 24 J. CLIN. ONCOL. 3020 (2006)

Petition for *Inter Partes* Review of USP 9,006,224

West-Ward Pharmaceuticals International Limited (“Petitioner” or “West-Ward”) requests *inter partes* review of claims 1-3 of United States Patent No. 9,006,224 (the “’224 patent”), titled “Neuroendocrine Tumor Treatment,” which according to USPTO records is assigned to Novartis AG (“Patent Owner” or “Novartis”). In *Par Pharmaceutical, Inc. v. Novartis AG*, IPR2016-01479 (Paper No. 8, February 15, 2017), the Board previously instituted an *inter partes* review on claims 1-3 of the ’224 patent on the same grounds (Grounds 1-4) upon which West-Ward relies below.

I. OVERVIEW

The Board should grant *inter partes* review because the ’224 patent claims nothing more than what was already well-known in the art. The ’224 patent claims methods of treating advanced pancreatic neuroendocrine tumors with a rapamycin derivative known as everolimus. The prior art, however, already taught treating these exact tumors with rapamycin and its derivatives. And everolimus was identified as having better bioavailability and presenting a “clinical advantage” over rapamycin. Further, everolimus specifically was taught to be effective in a recognized rat model of these pancreatic tumors. Additionally, unlike rapamycin, both everolimus and temsirolimus (another rapamycin derivative) had been shown

to be effective and well-tolerated in human cancer patients, and temsirolimus had been shown to be safe and effective in treating humans with advanced neuroendocrine tumors (“NETs”). Thus, it would have been obvious to use everolimus to treat advanced pancreatic NETs as recited in the claims.

II. REQUIREMENTS FOR PETITION FOR INTER PARTES REVIEW

A. Grounds for Standing (37 C.F.R. § 42.104(a))

Pursuant to 37 C.F.R. § 42.104(a), Petitioner hereby certifies that the '224 Patent is available for *inter partes* review. On June 10, 2015, Novartis filed a complaint in the District of Delaware alleging that Roxane, now West-Ward, infringed the '224 Patent. *Novartis Pharm. Corp. et al. v. Roxane Labs., Inc.*, No. 1:15-cv-474-RGA (D. Del.). On February 27, 2017, West-Ward was substituted as a party for Roxane in the District of Delaware litigation. Although more than one year has passed since Roxane, now West-Ward, was served with a complaint alleging infringement of the '224 Patent, Petitioner is not barred or estopped from *inter partes* review of the challenged claims of the '224 Patent on the grounds identified in this Petition. Neither Petitioner nor any privy of Petitioner has received a final written decision under 35 U.S.C. § 318(a) with respect to any claim of the '224 Patent on any ground that was raised or could have been raised

Petition for *Inter Partes* Review of USP 9,006,224

by Petitioner or its privies in any *inter partes* review, post grant review, or covered business method patent review. Moreover, the Petition (and an accompanying motion to join IPR2016-01479) is timely filed, *i.e.*, within one month of the February 15, 2017 institution of IPR2016-01479.

B. Notice of Lead and Backup Counsel and Service Information

Pursuant to 37 C.F.R. §§ 42.8(b)(3), 42.8(b)(4), and 42.10(a), West-Ward provides the following designation of Lead and Back-Up counsel.

LEAD COUNSEL	BACKUP COUNSEL
Keith A. Zullo (Reg. No. 37,975) kzullo@goodwinprocter.com Goodwin Procter LLP The New York Times Building 620 Eighth Avenue New York, NY 10018-1405 T: 212-813-8846; F: 646-558-4226	Marta E. Delsignore (Reg. No. 32,689) mdelsignore@goodwinprocter.com Goodwin Procter LLP The New York Times Building 620 Eighth Avenue New York, NY 10018-1405 T: 212-813-8822; F: 646-558-4079

A Power of Attorney is being filed concurrently herewith in accordance with 37 C.F.R. § 42.10(b). West-Ward consents to electronic service.

C. Notice of Real-Parties-in-Interest (37 C.F.R. § 42.8(b)(1))

Pursuant to 37 C.F.R. § 42.8(b)(1), Petitioner certifies that West-Ward is a real-party-in-interest for this proceeding. Out of an abundance of caution, and as a result of ongoing integration and reorganization activities, Petitioner identifies

the following additional entity as a real-party-in-interest who, going forward, may have control over this proceeding: Hikma Pharmaceuticals PLC and its U.S. subsidiaries Roxane Laboratories, Inc. and West-Ward Pharmaceuticals Corp. No other parties exercised or could have exercised control over this petition; no other parties funded or directed this petition. *See* Office Patent Trial Practice Guide, 77 Fed. Reg. 48759-60.

D. Notice of Related Matters (37 C.F.R. § 42.8(b)(2))

Pursuant to 37 C.F.R. § 42.8(b)(2), Petitioner is aware of the following District Court litigations: *Novartis Pharm. Corp. v. Breckenridge Pharm., Inc.*, C.A. No. 14-1043-RGA (D. Del.); *Novartis Pharm. Corp. v. Roxane Labs., Inc.*, C.A. No. 14-1196-RGA (D. Del.); *Novartis Pharm. Corp. v. Par Pharm., Inc.*, C.A. No. 14-1289-RGA (D. Del.); *Novartis Pharm. Corp. v. Par Pharm., Inc.*, C.A. No. 14-1494-RGA (D. Del.); *Novartis Pharm. Corp. v. Par Pharm., Inc.*, C.A. No. 15-78-RGA (D. Del.); *Novartis Pharm. Corp. v. Par Pharm., Inc.*, C.A. No. 15-475-RGA (D. Del.); *Novartis Pharm. Corp. v. Par Pharm., Inc.*, C.A. No. 15-1050-RGA (D. Del.); *Novartis Pharm. Corp. v. Roxane Labs., Inc.*, C.A. No. 14-1508-RGA (D. Del.); *Novartis Pharm. Corp. v. Roxane Labs., Inc.*, C.A. No. 15-128-RGA (D. Del.); and *Novartis Pharm. Corp. v. Breckenridge Pharm., Inc.*,

Petition for *Inter Partes* Review of USP 9,006,224

C.A. No. 16-431-RGA (D. Del.). Petitioner is also aware of the following petitions for *Inter Partes* Review of U.S. Patent No. 9,006,224: *Roxane Labs., Inc. v. Novartis AG*, IPR2016-01461; *Argentum Pharm. LLC v. Novartis AG*, IPR2017-01063; and *Par Pharm., Inc. v. Novartis AG*, IPR2016-01479.

Petitioner is also aware of the following petitions for *Inter Partes* Review of U.S. Patent No. 5,665,772: Nos. IPR2016-00084, -01023, -01059, -01102, and -01103.

Petitioner is also aware of the following petitions for *Inter Partes* Review: *Par Pharm., Inc. v. Novartis AG*, IPR2016-00074 (U.S. Patent No. 7,741,338); and *Par Pharm., Inc. v. Novartis AG*, IPR2016-00075 (U.S. Patent No. 7,297,703).

E. Fee for *Inter Partes* Review

The undersigned authorizes payment of \$23,000.00 for the fees specified by 37 C.F.R. § 42.15(a) for this Petition to be charged to Deposit Account No. 50-6989. The undersigned further authorizes payment for any additional fees that might be due in connection with this Petition to be charged to Deposit Account No. 50-6989.

F. Proof of Service

Proof of service of this petition on the Patent Owner at the correspondence address of record for the '224 patent is attached.

III. IDENTIFICATION OF CLAIMS BEING CHALLENGED (37 C.F.R. § 42.104(B))

For the reasons herein, the Board should find claims 1-3 unpatentable on the following grounds:

Ground 1. Claims 1-3 are unpatentable under 35 U.S.C. § 103 because they are rendered obvious by Öberg 2004 (Ex. 1027) in combination with Boulay 2004 (Ex. 1005) and O'Donnell (Ex. 1029).

Ground 2. Claim 2 is unpatentable under 35 U.S.C. § 103 because it is rendered obvious by Öberg 2004 (Ex. 1027) in combination with Boulay 2004 (Ex. 1005) and O'Donnell (Ex. 1029), in further view of Tabernero (Ex. 1038).

Ground 3. Claims 1-3 are unpatentable under 35 U.S.C. § 103 because they are rendered obvious by Boulay 2004 (Ex. 1005), O'Donnell (Ex. 1029), and Duran (Ex. 1011).

Ground 4. Claim 2 is unpatentable under 35 U.S.C. § 103 because it is rendered obvious by Boulay 2004 (Ex. 1005), O'Donnell (Ex. 1029), and Duran (Ex. 1011), in further view of Tabernero (Ex. 1038).

IV. SUMMARY OF THE ARGUMENT

The '224 patent claims methods of treating advanced pancreatic NETs by administering to a human subject in need thereof a therapeutically effective amount

of everolimus¹ after failure of cytotoxic chemotherapy. Ex. 1001, '224 patent at 26:65-27:8. Treating advanced pancreatic NETs (such as islet cell tumors) by administering a therapeutically effective amount (including 10 mg/day) of everolimus after cytotoxic treatment fails would have been obvious at the time of the purported invention, November 21, 2005.

First (*i.e.*, Grounds 1 and 2), Öberg 2004 disclosed rapamycin as a treatment for advanced pancreatic NETs after cytotoxic treatment failed, and one of skill would have understood that suggestion to include rapamycin's other known active derivatives that had been reported to be administered to human cancer patients, such as everolimus. Ex. 1027, Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶¶ 79, 83-92, 104. Everolimus was first disclosed in 1992, and subsequent preclinical and

¹ The claims of the '224 patent use the term 40-0-(2-hydroxyethyl)-rapamycin. This compound is also known in the art as everolimus, RAD001, SDZ RAD, and RAD. *E.g.*, Ex. 1001, '224 patent at 11:50-51; Ex. 1033, Rao at 621; Ex. 1003, Ratain Decl. ¶ 72. Sometimes Novartis and its predecessor Sandoz refer to 40-0-(2-hydroxyethyl)-rapamycin as Compound A. '224 patent at 11:66-67. For ease of reference, this Petition will primarily use the term "everolimus" in referencing this compound.

clinical research touted its activity and identifying it as having a “clinical advantage” over rapamycin. Ex. 1003, Ratain Decl. ¶¶ 72, 75-79. Further, everolimus would have been an obvious treatment choice because its efficacy in treating pancreatic NETs had been demonstrated in laboratory models and the prior art taught that everolimus was safe and effective in treating humans with solid tumors. Ex. 1005, Boulay 2004 at 254; Ex. 1029, O’Donnell at 803; Ex. 1003, Ratain Decl. ¶¶ 110-123. And although it would have been obvious to identify an appropriate dose, Tabernero explicitly taught using a unit dose of 10 mg/day of everolimus for treating solid tumors. Ex. 1038, Tabernero at 3007; Ex. 1003, Ratain Decl. ¶¶ 126-127, 152.

Second (*i.e.*, Grounds 3 and 4), Boulay 2004 demonstrated that everolimus was effective in treating pancreatic NETs in rats and would have suggested to one of ordinary skill to administer everolimus to humans with pancreatic NETs. A skilled artisan would have had a reasonable expectation that everolimus would be effective in pancreatic NETs because of the antitumor activity in this preclinical model. Ex. 1005, Boulay 2004 at 254; Ex. 1003, Ratain Decl. ¶ 112. Further, O’Donnell taught that everolimus was safe and effective for treating other tumors in humans. Ex. 1029, O’Donnell at 803; Ex. 1003, Ratain Decl. ¶¶ 119-123. One


of ordinary skill would have tried, and reasonably expected to succeed, using everolimus to treat advanced pancreatic NETs in humans after cytotoxic treatment failed because Duran had demonstrated that another well-known rapamycin derivative, temsirolimus,² was effective in treating advanced NETs. Ex. 1011, Duran at 3096; Ex. 1003, Ratain Decl. ¶¶ 129-131, 167.

To explain further, it was well-known as of November 2005 that rapamycin, an inhibitor of the protein mTOR, was a potent anti-tumor agent. Ex. 1003, Ratain Decl. ¶¶ 70-71, 83-92. Researchers had investigated the use of rapamycin in the treatment of a variety of cancers and tumor models, including two pancreatic cancer cell lines. *Id.* ¶¶ 70-71.


Because of rapamycin's promising results in that research, researchers investigated and identified rapamycin derivatives with similar anti-tumor and mTOR-inhibition properties, including everolimus and temsirolimus. *Id.* ¶¶ 72-82. As of November 2005, everolimus and temsirolimus were the two most studied rapamycin derivatives. *Id.* ¶ 75. Differing only at the C40 position (circled in

² Temsirolimus is also known as CCI-779 in the literature. Ex. 1033, Rao at 621; Ex. 1003, Ratain Decl. ¶ 73.

red), these two rapamycin derivatives have identical binding sites for their biological targets, mTOR and FKBP12³.



³ By November 2005, it was known that rapamycin and its derivatives first bind to the protein FKBP12 and then that rapamycin-FKBP12 complex interacts with mTOR to inhibit its activity. Ex. 1003, Ratain Decl. ¶ 83.



Id. ¶¶ 75, 91-92 (citing Ex. 1017, Huang 2002; Ex. 1039, Vignot at 528, Fig. 4). One of ordinary skill in November 2005 would have understood that both everolimus and temsirolimus have similar properties to each other and to rapamycin. Ex. 1003, Ratain Decl. ¶¶ 75-92.

Öberg 2004 suggested rapamycin (and therefore its known active derivatives) as a treatment for humans with advanced pancreatic NETs after the failure of chemotherapy. Ex. 1027, Öberg 2004 at 59; Ex. 1003, Ratain Decl. ¶¶ 100-104. Additionally, Boulay 2004 disclosed that everolimus was well-tolerated and effective in a rat model for pancreatic NETs that had been correlated to clinical efficacy in humans. Ex. 1005, Boulay 2004 at 254; Ex. 1003, Ratain

Decl. ¶¶ 110-117. Specifically, Boulay 2004 reported that administering everolimus as a monotherapy to rats injected with pancreatic NET tumor cells showed statistically significant antitumor activity, and was “well tolerated, with no significant body weight loss or mortalities observed.” Ex. 1005, Boulay 2004 at 254; Ex. 1003, Ratain Decl. ¶¶ 115-116. This model was reported to indicate likely clinical activity in PNET. Ex. 1003, Ratain Decl. ¶¶ 112.

A person of ordinary skill would have also known that both everolimus and temsirolimus are effective and well-tolerated in human cancer patients. Ex. 1029, O’Donnell at 803; Ex. 1003, Ratain Decl. ¶¶ 79-81; *see also* Ex. 1009, Dancey at 1105-1110; Ex. 1054, Dukart at 5:1-6:26; Ex. 1011, Duran at 3096. In particular, O’Donnell taught that everolimus exhibited anti-tumor effects and “was well tolerated with only mild degrees” of side effects. Ex. 1029, O’Donnell at 803; Ex. 1003, Ratain Decl. ¶ 120. And the prior art taught that everolimus was more bioavailable than rapamycin, with a “more favorable” pharmacokinetic profile, proving a “clinical advantage.” Ex. 1009, Dancey, at 1105-06; Ex. 1036, Schuler at 36-37, 41; Ex. 1003, Ratain Decl. ¶¶ 76-77. Further, Tabernero recommended that everolimus be administered in 10 mg daily doses as a monotherapy for treating advanced solid tumors. Ex. 1038, Tabernero at 3007.

In light of these teachings, a person of ordinary skill in the art seeking to treat patients with advanced pancreatic NETs after the failure of cytotoxic chemotherapy would have been motivated to combine the teachings of Öberg 2004, Boulay 2004, and O'Donnell to treat such tumors with everolimus as recited in the claims of the '224 patent. Ex. 1003, Ratain Decl. ¶¶ 132-138. Although one of ordinary skill would have been able to determine an appropriate dose using routine experimentation, a skilled artisan would have also incorporated the teaching of Tabernero that everolimus should be administered at 10 mg/day for the treatment of solid tumors, as recited in claim 2 of the '224 patent. Ex. 1038, Tabernero at 3007; Ex. 1003, Ratain Decl. ¶¶ 152-53, 159-160.

In addition, the prior art taught that everolimus had preclinical activity in a rat tumor model of pancreatic NETs, Boulay 2004 at 252-54, and clinical activity in treating humans with solid tumors, O'Donnell. A skilled artisan would have also been aware that the rapamycin derivative temsirolimus had been shown to have antitumor activity in humans with advanced neuroendocrine carcinomas, a subset of advanced NETs, who had previously been treated with cytotoxic chemotherapy. Ex. 1011, Duran at 3096; Ex. 1003, Ratain Decl. ¶¶ 94, 130.

Therefore, in seeking to treat humans with advanced pancreatic NETs, a skilled

artisan would have been motivated to combine the teachings of Boulay 2004, O'Donnell, and Duran, and in doing so, would have reasonably expected that everolimus would be effective in treating humans with advanced pancreatic NETs after failure with cytotoxic chemotherapy. Ex. 1003, Ratain Decl. ¶¶ 139-142, 161-174. And, as stated above, although a person of ordinary skill would have been able to determine an appropriate dose, including 10 mg/day, a skilled artisan would have also incorporated Taberero's teaching of a unit dose of 10 mg/day to treat solid tumors, which would include pancreatic NETs. Ex. 1038, Taberero at 3007; Ex. 1003, Ratain Decl. ¶¶ 125, 169-170, 176-177.

As described in more detail below, because the combinations of Öberg 2004, Boulay 2004, O'Donnell, and Taberero (*i.e.*, Grounds 1 and 2); and Boulay 2004, O'Donnell, Duran, and Taberero (*i.e.*, Grounds 3 and 4) each teach all elements of the claims, and are combinations motivated by a known problem in the field at the time of the alleged inventions, these combinations render claims 1-3 of the '224 patent obvious. *See In re Beattie*, 974 F.2d 1309, 1312 (Fed. Cir. 1992) ("As long as some motivation or suggestion to combine the references is provided by the prior art taken as a whole, the law does not require that the references be combined for the reasons contemplated by the inventor."); *see also KSR Int'l Co. v. Teleflex*

Inc., 550 U.S. 398, 421 (2007) (“When there is a design need or market pressure to solve a problem and there are a finite number of identified, predictable solutions, a person of ordinary skill has good reason to pursue the known options within his or her technical grasp.”).

V. OVERVIEW OF THE '224 PATENT

The '224 patent claims methods of using everolimus to treat humans with advanced pancreatic NETs after the failure of cytotoxic chemotherapy. Ex. 1001, '224 patent. Specifically, claim 1 recites:

1. A method for treating pancreatic neuroendocrine tumors, comprising administering to a human subject in need thereof a therapeutically effective amount of 40-0-(2-hydroxyethyl)-rapamycin as a monotherapy and wherein the tumors are advanced tumors after failure of cytotoxic chemotherapy.

Claim 2 recites a specific dose of everolimus:

2. The method of claim 1, wherein a unit dose of 40-0-(2-hydroxyethyl)-rapamycin is 10 mg/day.

And claim 3 recites a particular type of pancreatic NET:

3. The method of claim 1, wherein the tumor is [an] islet cell tumor.

Ex. 1001, '224 patent at 26:65-27:8.

The '224 patent describes the use of mTOR inhibitors, like rapamycin, everolimus, and temsirolimus, in the treatment of pancreatic NETs. *Id.* at 2:3-40. It further describes that rapamycin and its derivatives, such as everolimus and temsirolimus, inhibit mTOR activity through a complex with FKBP12. *Id.* at 1:6-15. And it states that mTOR inhibitors have potent antiproliferative properties, which makes them useful for cancer chemotherapy, and particularly for advanced solid tumors. *Id.* at 2:35-40.

The '224 patent does not include any preclinical data evidencing the effect of any mTOR inhibitor on pancreatic NETs. *See generally* Ex. 1001, '224 patent; Ex. 1003, Ratain Decl. ¶¶ 31-37. Instead, the '224 patent includes several prophetic examples, including administering everolimus (Compound A) to cancer cell lines to “show interesting antiproliferative activity in combination with another chemotherapeutic agent,” to determine “the immediate pharmacodynamic effect of the mTOR inhibitor,” to determine that “Compound A [everolimus] is able to restore activity of endocrine agents, like estrogen inhibitors and/or aromatase inhibitors” in breast cancer tumors. *Id.* at 25:49-26:27.

Nor does the '224 patent include any clinical data evidencing the effect of any mTOR inhibitor on pancreatic NETs in humans. *See generally* Ex. 1001, '224

patent; Ex. 1003, Ratain Decl. ¶ 37. Instead, the '224 patent includes several prophetic clinical trial protocols, including administering everolimus (Compound A) alone or in combination with Sandostatin LAR^{®4} in patients having carcinoid or islet cell cancer, evaluating the effect of adding everolimus (Compound A) to patients with advanced midgut carcinoid tumors already receiving Sandostatin LAR[®], and evaluating the effect of everolimus (Compound A) at a dosage of 10 mg/day in patients with measureable advanced pancreatic NETs (islet cell tumor) with or without Sandostatin LAR[®]. Ex. 1001, '224 patent at 26:28-64.

Boulay 2004 was submitted to the Patent Office during prosecution, but the Examiner never discussed or relied upon it. Ex. 1003, Ratain Decl. ¶ 49. Öberg 2004, O'Donnell, and Tabernero were neither submitted to the Patent

⁴ The active ingredient of Sandostain LAR[®] is octreotide acetate, a somatostatin analog. Ex. 1003, Ratain Decl. ¶ 35. Octreotide acetate was approved to treat symptoms of VIPomas, a type of pancreatic NET. *Id.* The use of somatostatin analogs as a treatment for pancreatic NETs was well-known. Ex. 1027, Öberg 2004 at 59-60.

Office nor considered by the Examiner during prosecution of the '224 patent.⁵

Id.

VI. THE PERSON OF ORDINARY SKILL IN THE ART

The '224 patent claims methods of treating pancreatic NETs in human patients by administering the known rapamycin derivative everolimus. Ex. 1003, Ratain Decl. ¶ 20; Ex. 1001, '224 patent. Accordingly, a person of ordinary skill

⁵ Novartis submitted Öberg 2004 and Tabernero to the Patent Office in the continuation application 14/608,644, on an IDS dated April 1, 2015, two weeks before the '224 patent issued, but did not submit Öberg 2004 or Tabernero to the Patent Office during prosecution of the parent '224 patent. Ex. 1056, Apr. 1, 2015 IDS at NPL refs. 8, 18. The Examiner cited Öberg 2004 in connection with the continuation application as teaching the use of somatostatin analogues as effective in inducing NET regression. Ex. 1057, Dec. 18, 2015, Office Action at 6-7. The Examiner has not cited or addressed Tabernero in connection with the continuation application. Thus, although these references have been before the Office as prior art in a continuation application, the Office has not considered arguments substantially similar to those presented in the instant Petition for the '224 patent.

in the art in November 2005 would have had (1) a medical degree (e.g., MD) with several years of specific experience in medical oncology, which generally includes board certification, as well as knowledge of oncology drug development and clinical pharmacology; or (2) a Ph.D. in cancer biology, molecular biology, medicinal chemistry, or a related field with several years of experience in oncology drug development and clinical pharmacology, including evaluating cancer therapeutics in *in vitro* and/or *in vivo* assays, as well as familiarity with the practice of medical oncology. Ex. 1003, Ratain Decl. ¶ 21. This description is approximate, and a higher level of education or skill might make up for less experience, and vice-versa. *Id.*

VII. CLAIM CONSTRUCTION

A. Applicable Law

In deciding whether to institute *inter partes* review, “[a] claim in an unexpired patent shall be given its broadest reasonable construction in light of the specification of the patent in which it appears.”⁶ 37 C.F.R. § 42.100(b); *Cuozzo*

⁶ The district court, in contrast, affords a claim term its “ordinary and customary meaning ... to a person of ordinary skill in the art in question at the time of the invention.” *Phillips v. AWH Corp.*, 415 F.3d 1303, 1313 (Fed. Cir.

Speed Techs., LLC v. Lee, 136 S.Ct. 2131,2142 (2016). This claim construction standard is different from — and broader than — that applied in district court.

Versata Dev. Grp., Inc. v. SAP Am., Inc., 793 F.3d 1306, 1327-28 (Fed. Cir. 2015).

Under the broadest reasonable interpretation, “claims should always be read in light of the specification and teachings in the underlying patent.” *Microsoft Corp.*

v. Proxyconn, Inc., 789 F.3d 1292, 1298 (Fed. Cir. 2015). The “PTO should also

consult the patent’s prosecution history in proceedings in which the patent has

been brought back to the agency for a second review.” *Id.* In addition, “[e]ven

under the broadest reasonable interpretation, the Board’s construction cannot be

divorced from the specification and the record evidence and must be consistent

with the one that those skilled in the art would reach.” *Id.* (quotation marks and

citations omitted).

To the extent there is any ambiguity regarding the “broadest reasonable construction” of a claim term, the ambiguity should be resolved in favor of the broader construction, absent amendment by the patent owner. Final Rules, 77 Fed. Reg. 48680, 48699 (Aug. 14, 2012) (“[T]he broader standard serves to identify ambiguities in the claims that can then be clarified through claim amendments.”).

2005) (en banc).

Consistent with the patent owner’s responsibility to clarify ambiguous terms, “the Office may take into consideration inconsistent statements made by a patent owner about claim scope, such as those submitted under 35 U.S.C. § 301(a), when applying the ‘broadest reasonable interpretation’ standard.” *Id.* Thus, while not controlling, claim constructions offered by a patent owner in related litigation (under the narrower standard applicable in district court) are relevant to the “broadest reasonable construction” applicable here. *See, e.g., SAP Am., Inc. v. Versata Dev. Grp., Inc.*, No. CBM2012-00001 (P.T.A.B. Jan. 9, 2013), Paper 36 at 8-9 (adopting petitioner’s proposed construction, in part, because it was “consistent with [p]atentee’s proposed construction in the related district court proceeding”).

B. Construction of Claim Terms

All claim terms—including those not specifically addressed in this section — have been accorded their broadest reasonable interpretation consistent with the specification of the ’224 patent as understood by one of ordinary skill in the art at the time of the alleged inventions. West-Ward provides the broadest reasonable interpretations of the following terms:

1. “pancreatic neuroendocrine tumor”

The term “pancreatic neuroendocrine tumor,” which is referred to as “pancreatic NET” in this petition, is recited in claim 1: “A method for treating pancreatic neuroendocrine tumors, comprising administering to a human subject ...” Ex. 1001, ’224 patent at claim 1.

A person of ordinary skill in the art at the time of the alleged inventions would understand that a NET is an abnormal growth of cells of the nervous or endocrine systems within or proximal to the pancreas. Ex. 1003, Ratain Decl. ¶ 53. The ’224 patent specification states that “Pancreatic neuroendocrine tumors as indicated herein e.g. include islet cell tumors, APUDomas, insulinomas, glucagonomas, nonfunctioning pancreatic NETS, pancreatic NETs associated with hypercalcemia, gastrinomas, VIPomas, somatostatinomas, GRFomas.” Ex. 1001, ’224 patent at 8:13-17. Reading this statement in the context of the entire patent, a skilled artisan would have understood the term “pancreatic neuroendocrine tumors” as used in the claims to include these tumors. Ex. 1003, Ratain Decl. ¶¶ 55-56.

Accordingly, the broadest reasonable construction of the claim term “pancreatic neuroendocrine tumor” is “abnormal growth of cells of the nervous or

endocrine systems in the pancreas, including, e.g., islet cell tumors, APUDomas, insulinomas, glucagonomas, nonfunctioning pancreatic NETS, pancreatic NETs associated with hypercalcemia, gastrinomas, VIPomas, somatostatinomas, and GRFomas.” Ex. 1003, Ratain Decl. ¶ 57.

2. “advanced tumors”

The term “advanced tumors” also appears in claim 1, “A method for treating pancreatic neuroendocrine tumors ... wherein the tumors are advanced tumors” Ex. 1001, ’224 patent at claim 1.

A person of ordinary skill in the art at the time of the alleged inventions would have understood the claim term “advanced tumors” to have its ordinary and customary meaning in the art that is consistent with the specification. As used by those of skill in the field of oncology, an “advanced” tumor is a tumor that is unresectable or metastatic. Ex. 1003, Ratain Decl. ¶ 59; *see also* Ex. 1023, Moertel at 520 (describing the patients with advanced islet cell carcinoma as having been identified with “proof of unresectable or metastatic islet-cell carcinoma.”).

This plain meaning is consistent with the ’224 patent specification, which correlates “advanced” tumors with “metastatic or unresectable” tumors. Ex. 1001,

'224 patent at 26:57-58 (“measurable advanced (metastatic or unresectable [*sic*, unresectable]) pancreatic neuroendocrine tumors”); Ex. 1003, Ratain Decl. ¶ 59.

An unresectable tumor is one that is unable to be completely removed by surgery. *Id.*

Accordingly, the broadest reasonable construction for “advanced tumors” is “tumors that are metastatic or unresectable.” *Id.* ¶ 60.

3. “unit dose”

The term “unit dose” appears in claim 2: “wherein a unit dose of [everolimus] is 10 mg/day.” Ex. 1001, '224 patent at 27:5-6.

A person of ordinary skill in the art at the time of the alleged inventions would have understood that the '224 patent uses the term “unit dose” consistent with its ordinary and customary meaning. Ex. 1003, Ratain Decl. ¶ 62. A person of ordinary skill would understand the term “unit dose” to refer to a single administration of the indicated dose. *Id.* The specification of the '224 patent does not use the term “unit dose,” however the specification does indicate that daily doses may be “administered in divided doses up to four times a day.” Ex. 1001, '224 patent at 10:27-36. The specification thus uses the term “divided dose[]” to

include the administration of a daily dose in multiple units. Ex. 1003, Ratain Decl.

¶ 62.

Accordingly, the broadest reasonable construction of the claim term “unit dose” is “dose administered as a single unit.” Ex. 1003, Ratain Decl. ¶ 63.

4. “islet cell tumor”

The term “islet cell tumor” appears in claim 3: “wherein the tumor is [an] islet cell tumor.” Ex. 1001, ’224 patent at 27:7-8.

A person of ordinary skill in the art at the time of the alleged inventions would have understood that the ’224 patent uses the term “islet cell tumor” consistent with its ordinary and customary meaning. Ex. 1003, Ratain Decl. ¶¶ 65-66. The ’224 patent specification indicates that islet cell tumors are equated with pancreatic NET, identifying islet cell tumors as a synonym: “pancreatic neuroendocrine tumors (islet cell tumor).” Ex. 1001, ’224 patent at 26:58-59.

Accordingly, the broadest reasonable construction of the claim term “islet cell tumor” is “abnormal growth of cells of the nervous or endocrine systems within the pancreas.” Ex. 1003, Ratain Decl. ¶ 67.

VIII. TECHNICAL BACKGROUND AND STATE OF THE ART AT THE TIME OF THE PURPORTED INVENTION


A. Rapamycin was well-known as a potent antitumor agent

Rapamycin was first discovered in a soil sample from Easter Island in the early 1970s, and scientists shortly identified its antifungal, antibiotic, and antitumor properties. Ex. 1003, Ratain Decl. ¶¶ 69-70; Ex. 1042, '171 patent at 2:7-4:14; Ex. 1013, Eng. Researchers have continued to study rapamycin in various cancer and tumor models. Ex. 1003, Ratain Decl. ¶¶ 70-71; *see also* Ex. 1044, '018 patent at 5:48-6:14 (describing the use of rapamycin in the treatment of skin carcinomas and malignant central nervous system carcinomas); Ex. 1015, Guba. At least as early as 1999, rapamycin was identified as an antitumor agent in two pancreatic cancer cell lines, MiaPaCa-2 and Panc-2. Ex. 1014, Grewe. Additionally, rapamycin was identified as a treatment for NETs — a category that includes pancreatic NETs — in 2004. Ex. 1003, Ratain Decl. ¶ 71; Ex. 1027, Öberg 2004 at 60.

B. Rapamycin derivatives, like everolimus and temsirolimus, were known to have similar biological activity to rapamycin

Rapamycin's promising activity sparked interest among researchers in identifying rapamycin derivatives — compounds similar to rapamycin, but with

small or minor modifications — with similar biological activity. *See, e.g.*, Ex. 1041, '803 patent; Ex. 1045, '036 patent; Ex. 1043, '883 patent; Ex. 1047, '730 patent at 1:47-3:6; Ex. 1049, '213 patent; Ex. 1003, Ratain Decl. ¶ 72. In 1992, Sandoz disclosed everolimus, a derivative formed by substituting the hydroxyl group at rapamycin's C40 position with a 2-hydroxyethyl group. Ex. 1048, '772 patent at 1:10-2:30; Ex. 1003, Ratain Decl. ¶ 72. That same year, American Home Products Corporation disclosed temsirolimus, a hydroxyester derivative of rapamycin also substituted at the C40 position. Ex. 1046, '718 patent; Ex. 1003, Ratain Decl. ¶¶ 73, 75. The chemical structures of rapamycin, everolimus, and temsirolimus are shown below, with the different groups at the C40 position circled in red:



Ex. 1003, Ratain Decl. ¶ 75.

As of November 2005, everolimus and temsirolimus were the most well-studied rapamycin derivatives, both clinically and in the laboratory. Ex. 1012, Dutcher; Ex. 1017, Huang 2002; Ex. 1039, Vignot; Ex. 1033, Rao; Ex. 1003, Ratain Decl. ¶ 75.

Contemporary publications touted the properties of everolimus to those of ordinary skill. Specifically, it was reported that everolimus has slightly lower pharmacological properties than rapamycin *in vitro*, but comparable properties *in vivo* and was developed to overcome formulation problems with rapamycin.

Ex. 1036, Schuler at 36-37, 38-40 (Tables 1-8), 41; Ex. 1003, Ratain Decl. ¶ 76.

One of ordinary skill would also have known that everolimus has slightly increased bioavailability and a shorter half-life than rapamycin. Ex. 1009, Dancey, at 1105-06; Ex. 1003, Ratain Decl. ¶ 77. Everolimus was reported to have “more favorable pharmacokinetic properties” which “provide a clinical advantage, i.e., it should be easier to handle and to monitor such a drug in clinical practice.” Ex. 1036, Schuler at 41; Ex. 1003, Ratain Decl. ¶ 76. Everolimus was also known to be well-tolerated with mild side effects when administered to human patients with solid tumors. Ex. 1029, O’Donnell at 803; Ex. 1003, Ratain Decl. ¶ 79.

Petition for *Inter Partes* Review of USP 9,006,224

By November 2005, everolimus was also well-known as having promising immunosuppressant and antitumor activities. For example, the prior art taught that everolimus “is a new, orally active rapamycin-derivative that is immunosuppressive and that efficiently prevents graft rejection in rat models ... and has therefore been selected for development.” Ex. 1036, Schuler at 36, Abstract; Ex. 1005, Boulay 2004 at 252 (“[Everolimus], an orally bioavailable derivative of rapamycin, demonstrates potent antiproliferative effects against a variety of mammalian cell types.... As a result of these properties, [everolimus] is being clinically developed both as an immunosuppressant ... and as a novel therapeutic in the fight against human cancer.”); Ex. 1003, Ratain Decl. ¶ 76. And Sandoz Ltd. described everolimus as having “particularly useful” properties for “proliferative disorders, e.g. tumors, hyper-proliferative skin disorder and the like.” Ex. 1048, ’772 patent at 3:22-4:10; *see also* Ex. 1036, Schuler at 36-37, 41 (describing selection of everolimus for clinical development and as having “more favorable” pharmacokinetic profile than rapamycin); Ex. 1003, Ratain Decl. ¶¶ 76-77. Other publications agreed that everolimus shows promising anti-tumor activity. Ex. 1029, O’Donnell; Ex. 1003, Ratain Decl. ¶¶ 78-79.

By November 2005, temsirolimus was also well-known as having promising antitumor activities. The prior art reported that temsirolimus shows “promising” results as an anticancer agent for advanced stage kidney cancer and “a phase III randomized trial[] is underway.” Ex. 1034, Sawyers at 344; Ex. 1003, Ratain Decl. ¶¶ 80-81. Temsirolimus was reported to be effective in reducing tumor mass when administered to mice engrafted with renal tumors, thus acting as an antineoplastic agent, “particularly for neoplasms which are refractory to standard therapy, or for whom standard therapy is not appropriate.” Ex. 1054, Dukart at 2:5-7, 4:10-25; Ex. 1003, Ratain Decl. ¶ 80. Two phase I clinical trials using temsirolimus had been conducted human patients with solid tumors and lymphomas. Ex. 1054, Dukart at 5:1-6:26. Tumor size was reported to be reduced in patients with a variety of cancers, including renal carcinoma, soft tissue carcinoma, breast cancer, neuroendocrine cancer of the lung, cervical cancer, uterine cancer, head and neck cancer, glioblastoma, non-small cell lung cancer, prostate cancer, pancreatic cancer, lymphoma, melanoma, small cell lung cancer, ovarian cancer, and colon cancer. *Id.* It was further known that temsirolimus showed “promising” results as an anticancer agent for advanced stage kidney cancer. Ex. 1034, Sawyers at 344; Ex. 1003, Ratain Decl. ¶ 81.


C. The mechanism of action for the immunosuppressant and antitumor activity of rapamycin and its derivatives was well-characterized

Before November 2005, researchers had made significant progress in identifying the mechanisms of action that made rapamycin and its derivatives — including everolimus and temsirolimus — biologically useful. It was known that rapamycin binds to the protein FKBP12 (FK-Binding Protein 12), and that this FKBP12-rapamycin complex inhibited the activity of the mTOR protein. Ex. 1037, Tolcher at S41-42; Ex. 1005, Boulay 2004 at 252 (“[Everolimus], like rapamycin, binds with high affinity to a ubiquitous intracellular receptor, the immunophilin FKBP12. This complex specifically interacts with ... mTOR ..., inhibiting downstream signaling events.”); Ex. 1003, Ratain Decl. ¶ 83.

It was also known that mTOR played a role in the Akt/PI3 kinase signal transduction pathway, which mediates proliferative signals and thus was “an attractive target for chemoprevention drug development.” Ex. 1037, Tolcher at S41-42; Ex. 1003, Ratain Decl. ¶ 84. Abnormalities in this Akt/PI3-mTOR pathway were known to be implicated in a number of different cancers, including pancreatic NETs. Ex. 1031, Perren at 1097, Abstract, 1101-02; Ex. 1009, Dancey at Table 1; Ex. 1003, Ratain Decl. ¶¶ 89, 96-97.

The FKBP12-rapamycin complex was known to inhibit the progression of the G1 phase of the cell cycle in osteosarcoma, liver, and T-cells, and interferes with mitogenic signaling pathways involved in G1 progression. Ex. 1006, Brown at 756; Ex. 1003, Ratain Decl. ¶ 85. This activity-the FKBP12- rapamycin complex inhibition of mTOR-was known to be responsible for the antiproliferative properties of rapamycin and its derivatives, including everolimus. Ex. 1039, Vignot at Abstract; Ex. 1033, Rao at 622; Ex. 1005, Boulay 2004 at 252-53; Ex. 1003, Ratain Decl. ¶ 86.

The mTOR and FKBP12-binding domains of rapamycin had been identified. *See, e.g.*, Ex. 1006, Brown; Ex. 1035, Schreiber at Fig. 5. Everolimus and temsirolimus were known to differ structurally from rapamycin only where circled in red below. This portion of the molecule is distanced from the mTOR and FKBP12-binding sites:



Ex. 1018, Huang 2003 at Fig. 1; Ex. 1033, Rao at Fig. 1; Ex. 1003, Ratain Decl.

¶ 91.

Thus, as of November 2005, a skilled artisan would have expected everolimus and temsirolimus each to have similar activity to their parent compound, rapamycin, in the mTOR signaling cascade. *See, e.g.*, Ex. 1018, Huang

2003 at Abstract; Ex. 1033, Rao at Abstract; Ex. 1003, Ratain Decl. ¶ 92. Further, a skilled artisan would have expected these mTOR inhibitors to be effective in tumor cells with hyperactivation of mTOR signaling, such as NETs. Ex. 1003, Ratain Decl. ¶¶ 88, 97-98; Ex. 1051, Wang 2002 at 140, Table 1, 144; Ex. 1028, Öberg & Ericksson at 755-56, Table 1.

IX. THE SCOPE AND CONTENT OF THE ASSERTED PRIOR ART

A. Öberg 2004 taught that humans with advanced pancreatic NETs should be treated with rapamycin as a monotherapy after cytotoxic therapy failed

Öberg 2004 (Ex. 1027) is prior art to the '224 patent under at least 35 U.S.C. § 102(b), because it published in April 2004, more than one year before the '224 patent's November 2005 priority date. *See* Öberg 2004; Ex. 1061, Bennett Decl. ¶¶ 44-53.

Öberg 2004 is a journal article disclosing the then-standard treatment protocols for NETs, a category that includes pancreatic NETs. *See, e.g.*, Öberg 2004 at 59; Ex. 1003, Ratain Decl. ¶¶ 100-108. Öberg 2004 reported the “clinical management of metastatic NE tumours.” *Id.* at 57. A person of skill in the art thus would understand that Öberg 2004's description of treating NETs includes the

treatment of advanced (*i.e.*, metastatic or unresectable) NETs. Ex. 1003, Ratain Decl. ¶¶ 101, 103.

Öberg 2004 graphically disclosed the therapy choices for treating NETs, including advanced NETs:




Fig. 1. Algorithm for the therapy of Neuroendocrine Tumours.

Öberg 2004 at Fig.1; Ex. 1003, Ratain Decl. ¶ 102. According to Öberg 2004, when treating NETs, surgery is the first option. Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶¶ 102-103. If the surgery is not successful or possible, then the tumor is unresectable and thus advanced. Ex. 1003, Ratain Decl. ¶ 59. For these advanced tumors, the next step is chosen based on whether the tumor is low-proliferative or high-proliferative. Öberg 2004 at

Fig. 1; Ex. 1003, Ratain Decl. ¶ 103. Low-proliferative tumors are treated first with biological therapies, such as interferon (IFN) or somatostatin inhibitors (SMS), and then cytotoxic therapy (*e.g.*, cytotoxic chemotherapeutics) if they fail to respond to biological therapies. Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶ 103. High-proliferative tumors are treated with cytotoxic therapy. Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶ 103. For any advanced (*i.e.*, unresectable) NETs which receive but are non-responsive to cytotoxic chemotherapies, experimental therapies are next recommended. Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶ 103. One of the disclosed experimental therapies is rapamycin. Öberg 2004 at Fig. 1, 60; Ex. 1003, Ratain Decl. ¶¶ 103-104.

Öberg 2004 expressly described rapamycin as “[a]nother interesting new compound” that “may block signal transduction through the m-TOR pathway. Clinical trials with this compound as a single agent or in combination with cytotoxic agents are planned.” Öberg 2004 at 60; Ex. 1003, Ratain Decl. ¶ 104. Öberg 2004 thus taught that rapamycin’s mTOR-inhibition properties are the reason for its use in treating NETs. Öberg 2004 at 60; Ratain Decl. ¶ 104. The recommendation to use rapamycin as a therapy for advanced NETs, including

advanced pancreatic NETs, was echoed in Wiedemann 2004, which identifies rapamycin as a “new targeted therap[y] that offer[s] new hope” “in order to improve current, rather limited treatment options especially in metastatic NET disease.” Ex. 1052, Wiedemann 2004 at 97; Ex. 1003, Ratain Decl. ¶¶ 105-108.

Öberg 2004 only differs from claims 1 and 3 of the '224 patent in that it does not explicitly disclose the use of everolimus and discloses rapamycin as a monotherapy in the treatment of human patients with advanced pancreatic NETs after the failure of cytotoxic chemotherapy. Öberg 2004 also does not include an explicit description of a 10 mg/day unit dose of everolimus as recited in claim 2 of the '224 patent.

B. Boulay 2004 taught that everolimus was well-tolerated and effective at treating pancreatic NETs in rat models

Boulay 2004 (Ex. 1005) is prior art to the '224 patent under at least 35 U.S.C. § 102(b), because it published in January 2004, more than one year before the '224 patent's November 2005 priority date. *See* Boulay 2004; Ex. 1061, Bennett Decl. ¶¶ 54-62.

Boulay 2004 disclosed that everolimus, “like rapamycin, binds with high affinity to ... FKBP12” and that “[t]his complex specifically interacts with ... mTOR” to “inhibit[] downstream signaling events.” *Id.* at 252. Boulay 2004

disclosed that everolimus was being clinically developed for anti-cancer therapeutics in humans. *Id.* (“[Everolimus], an orally bioavailable derivative of rapamycin, ... demonstrates potent antiproliferative effects against a variety of mammalian cell types As a result of these properties, [everolimus] is being clinically developed both as an immunosuppressant ... and as a novel therapeutic in the fight against human cancer.”); *see also id.* at 253; Ex. 1003, Ratain Decl. ¶ 110. Specifically, Boulay 2004 disclosed concurrent clinical trials of everolimus in humans with cancer. Boulay 2004 at 252, 260; Ex. 1003, Ratain Decl. ¶ 110.

Boulay 2004 also disclosed the use of everolimus as a monotherapy in treating rats with CA20948 tumors, which are a specific line of pancreatic NETs used in laboratory studies. Boulay 2004 at Abstract; Ex. 1003, Ratain Decl. ¶¶ 111-112. Boulay 2004 described this treatment as causing “antitumor activity characterized by statistically significant inhibition of tumor growth as compared with vehicle controls.” Boulay 2004 at 254; Ratain Decl. ¶ 113. Boulay 2004 disclosed that “[f]or all treatment schedules, [everolimus] was well tolerated” and elicited antitumor potency equivalent to that of the positive control, the cytotoxic agent 5-FU. Boulay 2004 at 254, 258; Ex. 1003, Ratain Decl. ¶¶ 115-116. Boulay 2004 disclosed that it is the “first full publication demonstrating significant

antitumor efficacy of a rapamycin derivative in an animal model of pancreatic cancer.” Boulay 2004 at 258; Ex. 1003, Ratain Decl. ¶ 117. Boulay 2004 disclosed that everolimus was “administered p.o. daily at 0.5 or 2.5 mg/kg (x6/week),” which indicates that the tumor-bearing rats in the study were given everolimus once per day by mouth at two different total doses (0.5 or 2.5 mg/kg) and that this once- daily treatment was administered six times over the course of a week. Ex. 1003, Ratain Decl. ¶ 114. Boulay 2004 further disclosed that everolimus was administered in doses of “0.5 mg/kg qd x6” and “2.5 mg/kg qd x6.” Boulay 2004 at 254, Fig. 1 at legend. One of ordinary skill would understand a dose administered “qd” is a dose administered once per day in a single administration. Ex. 1003, Ratain Decl. ¶ 114. Boulay 2004 thus disclosed the administration of everolimus as a monotherapy in a daily unit dose.

Boulay 2004 only differs from claims 1 and 3 of the '224 patent in that it does not (1) report efficacy in human patients with advanced pancreatic NETs (although it does disclose concurrent clinical trials in humans with cancer), or (2) explicitly identify administration after the failure of cytotoxic chemotherapy. Boulay 2004 also does not include an explicit description of a 10 mg/day unit dose

of everolimus as recited in claim 2 of the '224 patent, although it does disclose the daily administration of unit doses of everolimus.

C. O'Donnell taught that everolimus was well-tolerated and showed promise as an antitumor agent in human patients

O'Donnell (Ex. 1029) is prior art to the '224 patent under at least 35 U.S.C. § 102(b), because it published in June 2003, more than one year before the '224 patent's November 2005 priority date. *See* O'Donnell; Ex. 1061, Bennett Decl. ¶¶ 63-71.

O'Donnell disclosed the administration of various dosage levels of everolimus as a monotherapy to human patients with solid tumors. *Id.*; Ex. 1003, Ratain Decl. ¶ 119. O'Donnell disclosed that the treatments were “well tolerated” and only had “mild degrees” of side effects, and that “7/8 patients exhibited inhibition for at least 7 days.” O'Donnell at 803; Ex. 1003, Ratain Decl. ¶¶ 120-121. O'Donnell disclosed that patients were administered everolimus “once weekly” at 4 dose levels: 5, 10, 20, and 30 mg. O'Donnell at 803. A person of ordinary skill in the art would have understood from this that everolimus was administered in a single dose. Ex. 1003, Ratain Decl. ¶ 119. O'Donnell disclosed that other additional clinical studies had been initiated to explore everolimus's ability to treat human tumors. O'Donnell at 803; Ex. 1003, Ratain Decl. ¶ 123.

O'Donnell differs from claims 1 and 3 of the '224 patent only in that it does not (1) explicitly disclose the treatment of humans with advanced pancreatic NETs (although it does disclose clinical trials in humans with solid tumors), or (2) explicitly teach administration after the failure of cytotoxic chemotherapy. O'Donnell also does not include an explicit description of a 10 mg/day unit dose of everolimus as recited in claim 2 of the '224 patent, although it does disclose the administration of a unit dose of everolimus.

D. Tabernero taught that an appropriate dosage for humans taking everolimus for the treatment of advanced solid tumors was 10 mg/day

Tabernero (Ex. 1038) is prior art to the '224 patent under at least 35 U.S.C. § 102(a), because it was published in June 2005, before the '224 patent's November 2005 priority date. *See* Tabernero; Ex. 1061, Bennett Decl. ¶¶ 72-81.

Tabernero disclosed that everolimus was an mTOR-inhibitor, and further disclosed data from an investigation of everolimus's safety and recommended dosage levels for the treatment of cancer. *Id.* at 3007; Ex. 1003, Ratain Decl. ¶¶ 125-127. Tabernero disclosed that the safety and dosage levels were being investigated to develop everolimus further as an anti-tumor agent. Tabernero at 3007. Tabernero disclosed the treatment of human patients with advanced solid

tumors with everolimus in dosages of 20, 50, or 70 mg weekly, or 5 or 10 mg daily. Tabernero at 3007; Ex. 1003, Ratain Decl. ¶ 126. A person of ordinary skill in the art would have understood that in describing the doses as given “weekly” or “daily,” that the doses were given once per day or once per week. Ex. 1003, Ratain Decl. ¶ 126. Tabernero disclosed that everolimus was effective at inhibiting mTOR in the studied tumors. Tabernero at 3007; Ex. 1003, Ratain Decl. ¶ 125. Tabernero recommended a unit dose of 10 mg/day of everolimus as a monotherapy for further phase II/phase III clinical trials for treatment of solid tumors, which would include pancreatic NETs. Tabernero at 3007; Ex. 1003, Ratain Decl. ¶ 127.

E. Duran taught the use of temsirolimus in the treatment of human patients with advanced neuroendocrine carcinomas

Duran (Ex. 1011) is prior art to the '224 patent under at least 35 U.S.C. § 102(a), because it published in June 2005, before the '224 patent's November 2005 priority date. *See* Duran; Ex. 1061, Bennett Decl. ¶¶ 82-91.

Duran disclosed the use of the rapamycin-derivative temsirolimus in the treatment of human patients with metastatic neuroendocrine carcinomas (NECs), a subset of advanced NETs. *Id.* at 3096; Ex. 1003, Ratain Decl. ¶¶ 94, 129-130. Duran explicitly identified that NECs include “islet cell carcinomas,” a subset of islet cell tumors. Duran at 3096; Ex. 1003, Ratain Decl. ¶ 129. Of the twenty-

three patients who received temsirolimus treatment, eleven had previously undergone chemotherapy treatment. Duran; Ex. 1003, Ratain Decl. ¶ 130. Duran disclosed that temsirolimus “appears to have antitumor activity in NECs.” Duran at 3096; Ex. 1003, Ratain Decl. ¶ 130.

Duran differs from claims 1 and 3 of the ’224 patent in that it identifies the use of temsirolimus instead of everolimus in the treatment of patients with advanced NETs or islet cell tumors. Duran also does not include an explicit description of a 10 mg/day unit dose of everolimus as recited in claim 2 of the ’224 patent.

X. CLAIMS 1-3 WOULD HAVE BEEN OBVIOUS OVER THE PRIOR ART

For the specific reasons explained below, a person of ordinary skill in the art would have found claims 1-3 of the ’224 patent obvious in November 2005.

A. Legal Background

A patent claim is unpatentable if it is obvious in view of the prior art. 35 U.S.C. § 103. A finding of unpatentability requires that a claim be anticipated or obvious from the perspective of a skilled artisan at the time the invention was made. *Id.* In analyzing obviousness in light of the prior art, it is important to understand the scope of the claims, the level of skill in the relevant art, the scope

and content of the prior art, the differences between the prior art and the claims, and any secondary considerations of non-obviousness. *Graham v. John Deere Co.*, 383 U.S. 1 (1966).

“[I]f a technique has been used to improve one device, and a person of ordinary skill in the art would recognize that it would improve similar devices in the same way, using the technique is obvious unless its actual application is beyond his or her skill.” *KSR*, 550 U.S. at 417. There may also be a specific teaching, suggestion, or motivation to combine a prior art reference with another prior art reference. *E.g.*, *Ruiz v. A.B. Chance Co.*, 234 F.3d 654, 665 (Fed. Cir. 2000). Such a teaching, suggestion, or motivation to combine the prior art references may be explicit or implicit in the prior art. *Id.*

A patent is obvious when it “simply arranges old elements with each performing the same function it had been known to perform and yields no more than one would expect from such an arrangement,” as long as there is reason to combine the elements. *KSR*, 550 U.S. at 417-18 (internal quotation marks omitted). Similarly, “if a technique has been used to improve one device, and a person of ordinary skill in the art would recognize that it would improve similar

devices in the same way, using the technique is obvious unless its actual application is beyond his or her skill.” *Id.* at 417.

“[A]ny need or problem known in the field of endeavor at the time of invention and addressed by the patent can provide a reason for combining the elements in the manner claimed.” *KSR*, 550 U.S. at 420. “When there is a design need or market pressure to solve a problem and there are a finite number of identified, predictable solutions, a person of ordinary skill has good reason to pursue the known options within his or her technical grasp.” *Id.* at 421.

The obviousness inquiry takes “an expansive and flexible approach” to determine the scope and content of the prior art, differences between the prior art and the claims at issue, and the level of ordinary skill in the pertinent art. *Id.* at 407, 415. It considers “interrelated teachings of multiple patents; the effects of demands known to the design community or present in the marketplace; and the background knowledge possessed by a person having ordinary skill in the art, all in order to determine whether there was an apparent reason to combine the known elements in the fashion claimed by the patent at issue.” *Id.* at 418. “A person of ordinary skill is also a person of ordinary creativity, not an automaton.” *Id.* at 421. Thus a patent is obvious when it “simply arranges old elements with each

performing the same function it had been known to perform and yields no more than one would expect from such an arrangement,” as long as there is reason to combine the elements. *Id.* at 417-18. For instance, “[c]ombining two embodiments disclosed adjacent to each other in a prior art patent does not require a leap of inventiveness.” *Boston Sci. Scimed, Inc. v. Cordis Corp.*, 554 F.3d 982, 991 (Fed. Cir. 2009). Similarly, “if a technique has been used to improve one device, and a person of ordinary skill in the art would recognize that it would improve similar devices in the same way, using the technique is obvious unless its actual application is beyond his or her skill.” *KSR*, 550 U.S. at 417. The specific motivations for the combinations of the relied-upon references are discussed in the below section. The specific motivations to combine the references are discussed in the Grounds below.

B. Ground 1: Claims 1-3 would have been obvious in view of Öberg 2004, Boulay 2004, and O’Donnell

1. Claim 1

Claim 1 claims “[a] method for treating pancreatic neuroendocrine tumors, comprising administering to a human subject in need thereof a therapeutically effective amount of [everolimus] as a monotherapy and wherein the tumors are advanced tumors after failure of cytotoxic chemotherapy.” Ex. 1001, ’224 patent.

Petition for *Inter Partes* Review of USP 9,006,224

A person of ordinary skill in the art seeking to treat humans with advanced pancreatic NETs after the failure of cytotoxic chemotherapy would have looked to the teachings of Dr. Kjell Öberg. Before November 2005, Dr. Öberg was one of the preeminent clinical researchers on NETs. Ex. 1058, Öberg Biography; Ex. 1003, Ratain Decl. ¶ 132. He was one of the founders of the European Neuroendocrine Tumor Society (“ENETS”), whose primary aims was to “establish guidelines for the diagnosis and therapy of gastroenteropancreatic neuroendocrine tumors.” Ex. 1058, Öberg Biography; Ex. 1059, ENETS Info; Ex. 1003, Ratain Decl. ¶ 132. By November 2005, Dr. Öberg had also published numerous articles on NET treatment. Ex. 1058, Öberg Biography; Ex. 1003, Ratain Decl. ¶ 132.

Öberg 2004 taught the use of rapamycin as a monotherapy for the treatment of human patients with advanced pancreatic NETs after the failure of cytotoxic therapy. Ex. 1027, Öberg 2004; Ex. 1003, Ratain Decl. ¶ 133. Öberg 2004 depicted the then-standard treatment for NETs, which includes pancreatic NETs:




Fig. 1. Algorithm for the therapy of Neuroendocrine Tumours.

Ex. 1027, Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶¶ 133, 146. Öberg 2004 stated that NETs should be treated with cytotoxic therapy (*e.g.*, cytotoxic chemotherapeutics) for unresectable (*i.e.*, advanced) high proliferative tumors and unresectable (*i.e.*, advanced) low proliferative tumors that did not respond to biological therapies. Ex. 1027, Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶ 133. If cytotoxic therapies failed to treat any advanced NET, Öberg taught the use of various experimental therapies, including rapamycin. Ex. 1027, Öberg 2004 at Fig. 1; Ex. 1003, Ratain Decl. ¶¶ 133, 146.

Öberg described rapamycin as “[a]nother interesting new compound” that “may block signal transduction through the m-TOR pathway. Clinical trials with

this compound as a single agent or in combination with cytotoxic agents are planned.” Ex. 1027, Öberg 2004 at 60. A person of ordinary skill in the art would have understood from Öberg 2004 that rapamycin’s TOR-inhibition properties are the reason for its usefulness in treating advanced NETs. Ex. 1003, Ratain Decl. ¶ 146. Because Öberg 2004 taught that it is rapamycin’s ability to inhibit mTOR that made it a useful treatment for advanced NETs, a person of ordinary skill in the art would have understood that Öberg 2004’s teachings would extend to other rapamycin derivatives known to be mTOR inhibitors, such as everolimus and temsirolimus. Ex. 1003, Ratain Decl. ¶ 135. In fact, there were no reported data of rapamycin administered to cancer patients, in contrast to its known active derivatives, everolimus and temsirolimus. *Id.* ¶¶ 104, 135. As such, a skilled artisan would have understood Öberg 2004’s reference to rapamycin to include the derivatives with known clinical safety and efficacy in cancer patients. *Id.*

By November 2005, everolimus and temsirolimus were the most studied rapamycin derivatives for their clinical efficacy in various cancers. Ex. 1029, O’Donnell; Ex. 1054, Dukart; Ex. 1009, Dancey; Ex. 1038, Tabernero; Ex. 1011, Duran; Ex. 1003, Ratain Decl. ¶¶ 75, 135-136. And everolimus was known to have a “clinical advantage” over rapamycin and be more bioavailable. Ex. 1003,

Ratain Decl. ¶¶ 76-77. A person of ordinary skill in the art seeking to treat human patients with advanced pancreatic NETs after the failure of cytotoxic chemotherapy using the “experimental drug” specifically suggested by Öberg 2004 would have also looked to rapamycin derivatives with similar antitumor and mTOR-inhibition properties to rapamycin, especially those that had demonstrated safety and efficacy in human cancer patients or those shown to be effective in treating pancreatic NETs in preclinical models. Ex. 1003, Ratain Decl. ¶¶ 137-138, 149-151.

Specifically, a skilled artisan would have looked to other rapamycin derivatives to use as a treatment for pancreatic NETs as taught in Öberg 2004, and Boulay 2004 reference disclosed data showing that the use of everolimus in a rat pancreatic NET model was effective and well-tolerated as a monotherapy. Ex. 1005, Boulay 2004; Ex. 1003, Ratain Decl. ¶¶ 111-113, 137.

Boulay 2004 disclosed the successful use of everolimus as a monotherapy in a rat model of pancreatic NETs. Ex. 1005, Boulay 2004 at 252-54; Ex. 1003, Ratain Decl. ¶¶ 111-113. Specifically, rats with CA20948 pancreatic NETs were given daily doses of 0.5 or 2.5 mg/kg of everolimus. Ex. 1005, Boulay 2004 at 252. The treatments were “well tolerated, with no significant body weight loss or

mortalities observed,” and resulted in statistically significant antitumor activity.

Id. at 254.

A skilled artisan would have been motivated by Boulay 2004 to substitute rapamycin with everolimus for the treatment of humans with pancreatic NETs as taught in Öberg 2004 because of the encouraging data on its effectiveness in this rat model of pancreatic NET. Ex. 1003, Ratain Decl. ¶¶ 112, 137. And a skilled artisan would have had a reasonable expectation that everolimus would be effective in humans because O’Donnell taught that the use of everolimus in human cancer patients was effective and safe. Ex. 1029, O’Donnell; Ex. 1003, Ratain Decl. ¶¶ 120-121, 138. O’Donnell disclosed the use of various dosage levels of everolimus in the treatment of human patients with solid tumors and taught that the treatments were “well tolerated” and only had mild side effects. Ex. 1029, O’Donnell; Ex. 1003, Ratain Decl. ¶¶ 120-121, 138. O’Donnell disclosed that other additional clinical studies had been initiated to explore everolimus’s ability to treat human tumors. Ex. 1029, O’Donnell; Ex. 1003, Ratain Decl. ¶ 123.

As discussed, a skilled artisan would have been motivated to combine the teachings of Öberg 2004, Boulay 2004, and O’Donnell, and this combination “simply arranges old elements with each performing the same function it had been

known to perform and yields no more than one would expect from such an arrangement.” *KSR*, 550 U.S. at 417, 420 (“[A]ny need or problem known in the field of endeavor at the time of invention and addressed by the patent can provide a reason for combining the elements in the manner claimed.”); *see also Beattie*, 974 F.2d at 1312.

Here, cancer patients with advanced pancreatic NETs need additional treatment after chemotherapy fails. Ex. 1003, Ratain Decl. ¶ 99. Öberg 2004 taught using rapamycin as a monotherapy to treat advanced pancreatic NETs after the failure of cytotoxic chemotherapy; Boulay 2004 taught that everolimus was effective as a monotherapy in treating pancreatic NETs in mammals; and O’Donnell taught that everolimus was safe and effective in humans for the treatment of solid tumors. A person of ordinary skill in the art would have had a reasonable expectation of success in using everolimus as a monotherapy instead of rapamycin as taught in Öberg 2004 because Boulay 2004 demonstrated that everolimus was effective in treating pancreatic NETs in mammals. If everolimus was effective in the preclinical model for pancreatic NETs, a skilled artisan would reasonably have expected that it would be effective in treating humans with

pancreatic NETs. Ex. 1003, Ratain Decl. ¶ 112. Accordingly, claim 1 would have been obvious over this combination. *Id.* ¶¶ 149-151.

2. Claim 2

Claim 2 depends from claim 1 and further recites “wherein a unit dose of [everolimus] is 10 mg/day.” The limitations of claim 2 are obvious for all the reasons discussed above for claim 1.

O’Donnell disclosed that everolimus was administered as a unit dose weekly at 5, 10, 20, and 30 mg for treatment of solid tumors, but did not explicitly disclose a unit dose of 10 mg/day. Ex. 1029, O’Donnell. Boulay 2004 further disclosed that everolimus in a unit dose of 0.5 or 2.5 mg/kg daily resulted in antitumor activity in the rat pancreatic NET model, but it too did not explicitly disclose a unit dose to a human of 10 mg/day. Ex. 1005, Boulay 2004 at 253, 254. Other reports indicated that everolimus was administered at 5-10 mg/day in clinical studies. Ex. 1039, Vignot at Table 1; Ex. 1003, Ratain Decl. ¶ 152.

Regardless, the dose recited in claim 2 would have been obvious because the ’224 patent does not set forth any human clinical or laboratory data showing that a daily unit dose of 10 mg is optimal compared to any other dose of everolimus. *Merck & Co. v. Teva Pharm. USA, Inc.*, 395 F.3d 1364, 1374 (Fed. Cir. 2005)

(finding claim limitation regarding dosage obvious when the patent “sets forth no human clinical or laboratory data showing the safety and tolerability of the treatment methods claimed by the patent” to distinguish any “concerns for dose-related GI problems”); Ex. 1003, Ratain Decl. ¶ 153; *see generally* Ex. 1001, ’224 patent.

Further, identifying an appropriate and effective dose for the treatment of tumors (*e.g.*, dose titration) is a standard exercise requiring nothing more than routine experimentation. Ex. 1003, Ratain Decl. ¶ 153. Identifying such a dose “flows from the ‘normal desire of scientists or artisans to improve upon what is already generally known.’” *Pfizer, Inc. v. Apotex, Inc.*, 480 F.3d 1348, 1368 (Fed. Cir. 2007), quoting *In re Peterson*, 315 F.3d 1325, 1330 (Fed. Cir. 2003). Thus, a person of ordinary skill in the would have been motivated to identify an appropriate dose of everolimus to effectively treat humans with advanced pancreatic NETs, including a unit dose of 10 mg/day. Ex. 1003, Ratain Decl. ¶ 153. As such, claim 2 would have been obvious over this combination.

3. Claim 3

Claim 3 depends from claim 1 and further recites “wherein the tumor is [an] islet cell tumor.” The limitations of claim 1 are obvious for all the reasons discussed above.

With respect to claim 3, an islet cell tumor is a subset of pancreatic NETs and thus teachings in the art directed to the treatment of NETs and pancreatic NETs would apply equally to islet cell tumors. Ex. 1003, Ratain Decl. ¶¶ 156-157. As such, claim 3 would have been obvious over this combination.

C. Ground 2: Claim 2 would have been obvious in view of Öberg 2004, Boulay 2004, O’Donnell, and Tabernero

Claim 2 depends from claim 1 and further recites “wherein a unit dose of [everolimus] is 10 mg/day,” and would have been obvious over the above-discussed Öberg 2004 combination, in further view of Tabernero.

The only limitation present in claim 2 but not in claim 1, is the 10 mg/day unit dose limitation. Ex. 1001, ’224 patent. Although dose titration to identify effective doses is a routine skill known to a skilled artisan, a skilled artisan would have also searched for available everolimus dosage information. Ex. 1003, Ratain Decl. ¶¶ 143, 152, 159. Tabernero expressly recommended a unit dose of 10 mg/day of everolimus for treating humans with advanced solid tumors. Ex. 1038,

Tabernero. Specifically, after examining the safety and proper dosage levels of everolimus as an antitumor agent for advanced solid tumors, Tabernero disclosed that a unit dose of 10 mg/day was well-tolerated and effective and recommended that dosage for further clinical trials with everolimus as a single agent (*i.e.*, as a monotherapy). Ex. 1038, Tabernero; Ex. 1003, Ratain Decl. ¶ 159.

Tabernero expressly identified a recommended daily unit dose for treating advanced solid tumors and taught that this unit dose was well-tolerated and effective. A skilled artisan would therefore have had a reasonable expectation of success in achieving an effective therapy for advanced pancreatic NETs after failure of cytotoxic chemotherapy in administering a unit dose of 10 mg/day of everolimus in combining Öberg 2004, Boulay 2004, and O'Donnell with Tabernero. Ex. 1003, Ratain Decl. ¶¶ 143, 160. Accordingly, claim 2 is obvious over this combination.

D. Ground 3: Claims 1-3 would have been obvious in view of Boulay 2004, O'Donnell, and Duran

1. Claim 1

A skilled artisan seeking to treat advanced pancreatic NETs in humans after the failure of cytotoxic chemotherapy would have identified Boulay 2004, which disclosed that everolimus treatment was successful as a monotherapy in a rat

pancreatic NET model. Ex. 1005, Boulay 2004; Ex. 1003, Ratain Decl. ¶¶ 112, 141. A skilled artisan would have known that everolimus would be safe and effective in human cancer patients. Ex. 1003, Ratain Decl. ¶¶ 119-121. O'Donnell taught that everolimus treatment was well-tolerated and effective in treating human cancer patients with solid tumors. Ex. 1029, O'Donnell; Ex. 1003, Ratain Decl. ¶¶ 119-123. Duran further taught that the related mTOR inhibitor and similar rapamycin derivative, temsirolimus, had been shown to be safe and effective as a monotherapy in treating human patients with advanced NET previously treated with cytotoxic chemotherapy. Ex. 1011, Duran; Ex. 1003, Ratain Decl. ¶¶ 129-131, 142. Because temsirolimus was known to have similar anti-tumor properties as everolimus and was known to be an mTOR inhibitor like both rapamycin and everolimus and was known to be effective in humans with NETs, a skilled artisan would have a reasonable expectation that everolimus would also be safe and effective as a monotherapy in treating human patients with advanced NET previously treated with chemotherapy. Ex. 1003, Ratain Decl. ¶¶ 75, 82, 87-88, 91- 92, 142. Thus, the prior art taught that everolimus was effective as a monotherapy in a rat model of pancreatic NET (Boulay 2004), everolimus was safe and effective as a monotherapy in treating solid tumors in human patients

(O'Donnell), and the related rapamycin derivative temsirolimus was safe and effective as a monotherapy in treating advanced NET in humans after failure of chemotherapy (Duran). A skilled artisan would have been motivated to administer everolimus after failure of cytotoxic chemotherapy because such therapy was the “gold standard” but associated with low success rates. Ex. 1003, Ratain Decl. ¶ 166. As such, a skilled artisan would have been aware of the need for additional treatment options after chemotherapy failed. *Id.* ¶¶ 99, 166.

A skilled artisan would have had a reasonable expectation of success in administering everolimus as a monotherapy to human patients with advanced pancreatic NETs after the failure of cytotoxic chemotherapy in light of the teachings of Boulay 2004, O'Donnell, and Duran. Ex. 1003, Ratain Decl. ¶ 167. Accordingly, claim 1 of the '224 patent would have been obvious in view of this combination.

2. Claim 2

Claim 2 depends from claim 1 and further recites “wherein a unit dose of [everolimus] is 10 mg/day.” The limitations of claim 2 are obvious for all the reasons discussed above for claim 1.

O'Donnell disclosed that everolimus was administered to humans in weekly unit doses at 5, 10, 20, and 30 mg for treatment of solid tumors but did not explicitly disclose a unit dose of 10 mg/day. Ex. 1029, O'Donnell. Boulay 2004 further disclosed that everolimus administered in daily unit doses of 0.5 or 2.5 mg/kg resulted in antitumor activity in the rat pancreatic NET but did not explicitly disclose a unit dose to a human of 10 mg/day. Ex. 1005, Boulay 2004 at 253, 254. Other clinical reports disclosed everolimus administered at 5-10 mg/day. Ex. 1039, Vignot at Table 1; Ex. 1003, Ratain Decl. ¶ 169.

Regardless, the dose recited in claim 2 would have been obvious because the '224 patent does not set forth any human clinical or laboratory data showing that a daily unit dose of 10 mg is optimal compared to any other dose of everolimus. *Merck*, 395 F.3d at 1374 (finding claim limitation regarding dosage obvious when the patent “sets forth no human clinical or laboratory data showing the safety and tolerability of the treatment methods claimed by the patent” to distinguish any “concerns for dose-related GI problems”); Ex. 1003, Ratain Decl. ¶ 170; *see generally* Ex. 1001, '224 patent.

Further, identifying an appropriate and effective dose for the treatment of tumors (*e.g.*, dose titration) is a standard exercise requiring nothing more than

routine experimentation. Ex. 1003, Ratain Decl. ¶ 169. Identifying such a dose “flows from the ‘normal desire of scientists or artisans to improve upon what is already generally known.’” *Pfizer*, 480 F.3d at 1368, quoting *Peterson*, 315 F.3d at 1330. Thus, a person of ordinary skill in the art would have been motivated to identify an appropriate dose of everolimus to effectively treat humans with advanced pancreatic NETs, including a unit dose of 10 mg/day. As such, claim 2 would have been obvious over this combination.

3. Claim 3

Claim 3 depends from claim 1 and further recites “wherein the tumor is [an] islet cell tumor.” The limitations of claim 1 are obvious for all the reasons discussed above.

With respect to claim 3, an islet cell tumor is a subset of pancreatic NETs and thus teachings in the art directed to the treatment of NETs and pancreatic NETs would apply equally to islet cell tumors. Ex. 1003, Ratain Decl. ¶ 173. Indeed, Duran explicitly identified islet cell carcinomas as included within NEC. Ex. 1011, Duran at 3096. As such, claim 3 would have been obvious over this combination.

E. Ground 4: Claim 2 of the '224 patent is invalid as obvious in view of Boulay 2004, O'Donnell, Duran, and Tabernero

Claim 2 depends from claim 1 and further recites “wherein a unit dose of [everolimus] is 10 mg/day,” and would have been obvious over the above-discussed Boulay 2004 combination, in further view of Tabernero.

The only limitation present in claim 2 but not in claim 1, is the 10 mg/day unit dose limitation. Ex. 1001, '224 patent. Although dose titration to identify effective doses is a routine skill known to a skilled artisan, a skilled artisan would also search for available dosage information. Ex. 1003, Ratain Decl. ¶¶ 143, 169, 176. Tabernero expressly recommended a unit dose of 10 mg/day of everolimus for treating human tumors. Ex. 1038, Tabernero. Specifically, after examining the safety and proper dosage levels of everolimus as an antitumor agent, Tabernero disclosed that a unit dose of 10 mg/day was well-tolerated and effective and recommended that dosage for further clinical trials with everolimus as a single agent (*i.e.*, as a monotherapy). Ex. 1038, Tabernero; Ex. 1003, Ratain Decl. ¶ 176.

Tabernero expressly identified a recommended daily unit dose for treating advanced solid tumors and taught that it was well-tolerated and effective. A skilled artisan would therefore have had a reasonable expectation of success in achieving an effective therapy for advanced pancreatic NETs after failure of

cytotoxic chemotherapy in administering a unit dose of 10 mg/day in combining the teachings of Boulay 2004, O'Donnell, and Duran with Taberbero. Ex. 1003, Ratain Decl. ¶¶ 143, 177. Accordingly, claim 2 is obvious over this combination.

XI. SECONDARY CONSIDERATIONS FAIL TO OVERCOME THE STRONG EVIDENCE OF OBVIOUSNESS

To overcome West-Ward's strong *prima facie* obviousness showing set forth in the four proposed grounds above, Novartis may attempt to come forward with secondary considerations of nonobviousness. Secondary considerations can include (i) long-felt need, (ii) unexpected results, (iii) skepticism of the invention, (iv) teaching away from the invention, and (vii) coping by others. *Graham*, 383 U.S. at 15-17. West-Ward is unaware of any secondary considerations that would support a finding of nonobviousness for claims 1-3. Ex. 1003, Ratain Decl. ¶ 178. Moreover, even where relevant secondary considerations are present, they may not make claims nonobvious, particularly where — like here — a strong *prima facie* case exists. *See, e.g., Leapfrog Enters., Inc. v. Fisher-Price, Inc.*, 485 F.3d 1157, 1162 (Fed. Cir. 2007).

Novartis may allege that the method claims of the '224 patent satisfied a long-felt but unmet need for treating pancreatic NETs. But other treatments were available as of November 2005 for the treatment of pancreatic NETs, as identified

in Öberg 2004. Ex. 1027, Öberg 2004 at 57-60. Thus, “others had previously solved” any “long-felt need” allegedly met by the ’224 patent. *In re PepperBall Techs., Inc.*, 469 F. App’x 878, 882-83 (Fed. Cir. 2012).

Further, sales of everolimus for the treatment of pancreatic NETs do not support any objective indicia of the non-obviousness of the ’224 patent claims. The ’224 patent claims require administering everolimus, a compound whose use remains subject to separate patent coverage even today. Ex. 1048, ’772 patent at claim 10. As a result, “no entity other than [Novartis] could have successfully brought [the claimed methods] to market.” *Galderma Labs., L.P. v. Tolmar, Inc.*, 737 F.3d 731, 740 (Fed. Cir. 2013). Because of this barrier to market entry, any argument that commercial sales of Novartis’s Affinitor[®] (everolimus) product evidence non-obviousness is weak at best, especially in view of the strong *prima facie* case discussed above. *Merck*, 395 F.3d at 1376-77; *SIBIA Neurosciences, Inc. v. Cadus Pharm. Corp.*, 225 F.3d 1349, 1358 (Fed. Cir. 2000).

Finally, if Novartis alleges that copying by ANDA applicants supports nonobviousness, such an argument has been dismissed as “not probative of nonobviousness” in the ANDA context. *Bayer Healthcare Pharm., Inc. v. Watson Pharm., Inc.*, 713 F.3d 1369, 1377 (Fed. Cir. 2013).

CERTIFICATE OF SERVICE

The undersigned certifies that a complete copy of this Petition for *Inter Partes* Review of U.S. Patent No. 9,006,224 and all Exhibits and other documents filed together with this Petition were served on this 13th day of March, 2017 on the Patent Owner by serving via overnight delivery to the official correspondence address of record for U.S. Patent No. 9,006,224 shown in PAIR and Novartis Pharmaceutical Corporation's current litigation counsel:

Novartis Pharmaceutical Corporation
Intellectual Property Department
Attn: Peter J. Waibel, Esq. (Head of Patent Litigation)
One Health Plaza 433/2
East Hanover, NJ 07936-1080

Nicholas Kallas
Fitzpatrick, Cella, Harper & Scinto
1290 Avenue of the Americas
New York, NY 10104-3800

In addition, counsel for Novartis AG was served via FedEx International

Priority at the following address:

Novartis International AG
Novartis Campus
Attn: Alisa A. Harbin, Esq. (Head, Group Litigation and
Intellectual property)
Forum 1-1.20
Basel, CH-4002
Switzerland

Courtesy copies of the foregoing were also served via overnight delivery on the counsel of record for the Petitioner and Patent Owner in *Par Pharmaceuticals, Inc. v. Novartis AG*, IPR2016-01479 as follows:

Par Pharmaceutical, Inc.

Daniel Brown
Latham & Watkins LLP
885 Third Avenue
New York, NY 10022-4834

Jonathan M. Strang
Latham & Watkins LLP
555 Eleventh Street, NW, Ste. 1000
Washington, DC 20004-1304

Novartis AG

Nicholas N. Kallas
Fitzpatrick, Cella, Harper & Scinto
1290 Avenue of the Americas
New York, NY 10104-3800

March 13, 2017

/Keith A. Zullo/
Keith A. Zullo (Reg. No. 37,975)
GOODWIN PROCTER LLP
The New York Times Building
620 Eighth Avenue
New York, NY 10018-1405
Tel: 212-813-8846
Fax: 646-558-4226
kzullo@goodwinprocter.com

*Counsel for Petitioner West-Ward
Pharmaceuticals International Limited*

CERTIFICATE OF COMPLIANCE WITH 37 C.F.R. § 42.24

I hereby certify that this Petition complies with the word count limitation of 37 C.F.R. § 42.24(a)(1)(i) because the Petition contains 11,469 words, excluding the cover page, signature block, and the parts of the Petition exempted by 37 C.F.R. § 42.24(a)(1).

March 13, 2017

/Keith A. Zullo/
Keith A. Zullo (Reg. No. 37,975)
GOODWIN PROCTER LLP
The New York Times Building
620 Eighth Avenue
New York, NY 10018-1405
Tel: 212-813-8846
Fax: 646-558-4226
kzullo@goodwinprocter.com

*Counsel for Petitioner West-Ward
Pharmaceuticals International Limited*